

DEHRADUN PUBLIC SCHOOL
ASSIGNMENT (2022-23)
SUBJECT- MATHEMATICS (041)
CLASS - X


CHAPTER - 1 (REAL NUMBERS)

Solve the following questions:

Q1. The HCF of 2472, 1284 and a third number N is 12. If their LCM is $2^3 \times 3^2 \times 5 \times 103 \times 107$, then the number N is:

- a. $2^3 \times 3^2 \times 7$ b. $2^2 \times 3^3 \times 103$ c. $2^2 \times 3^2 \times 5$ d. $2^4 \times 3^2 \times 11$

Q2. The values of x and y in the given figure are:



- a. 7, 13 b. 13, 7 c. 9, 12 d. 12, 9

Q3. HCF of two numbers is 23 and their LCM is 1449. If one of the numbers is 161, then the other number is:

- a. 207 b. 307 c. 1449 d. None of these

Q4. If p and q are positive integers such that $p = ab^2$ and $q = a^3 b$ where a, b are prime numbers, then HCF (p, q) is:

- a. ab b. a^2b^2 c. a^3b^2 d. a^3b^3

Q5. In a school, there are two sections - section A and section B of class X. There are 32 students in section A and 36 students in section B. Determine the minimum number of books required for their class library so that they can be distributed equally among students of section A or section B :

- a. 288 b. 388 c. 208 d. None of these

Q6. The product of a non-zero rational and an irrational number is _____.

- a. always irrational b. always rational
c. rational or irrational d. none of these

Q7. The least number that is divisible by all the numbers from 1 to 10 (both inclusive) is:

- a. 10 b. 100 c. 504 d. 2520

Q8. The HCF and the LCM of 12, 21, 15 respectively are:

- a. 3, 140 b. 12, 420 c. 3, 420 d. 420, 2

- Q9.** The exponent of 2 in the prime factorization of 144 is _____.
- a. 2 b. 4 c. 1 d. 6
- Q10.** If sum of two numbers is 1215 and their HCF is 81, then the possible number of pairs of such numbers are:
- a. 2 b. 3 c. 4 d. 5
- Q11.** If $xy = 180$ and $\text{HCF}(x, y) = 3$, then find the LCM (x, y) .
- Q12.** Find the least positive integer divisible by 20 and 24.
- Q13.** If $\text{HCF}(a, b) = 12$ and $a \times b = 3600$, then find LCM (a, b) .
- Q14.** If p is a prime number and p divides a^2 (where 'a' is a positive integer), then p divides a . Is it true?
- Q15.** Is it true to say that the product of a rational number and an irrational number may be a rational or an irrational?
- Q16.** Find the HCF and LCM of 404 and 96. Verify that $\text{HCF} \times \text{LCM} = \text{Product of the two numbers}$.
- Q17.** 3 bells ring at an interval of 4, 7 and 14 minutes. All three bells rang at 6 am, when the three balls will the ring together next?
- Q18.** Prove $\sqrt{5}$ is irrational.
- Q19.** Prove that $4 - 2\sqrt{5}$ is an irrational.
- Q20.** $\text{HCF}(306, 657) = 9$, find LCM $(306, 657)$.
- Q21.** Explain why $17 \times 11 \times 13 + 11$ is a composite number.
- Q22.** Find the LCM and HCF of 15, 18 and 45 by the prime factorization method.
- Q23.** Mathematics teacher of a school decided to have maximum number of mixed sections for a team. Each section has to accommodate equal number of boys and equal number of girls. What is the number of such sections if there are 372 boys and 444 girls?
- Q24.** Prove that $\sqrt{2}$ is an irrational number. Hence, show that $5 + 3\sqrt{2}$ is also an irrational number.

Case-study based questions:

- Q25.** To enhance the reading skills of grade X students, the school nominates you and two of your friends to set up a class library. There are two sections- section A and section B of grade X. There are 32 students in section A and 36 students in section B.



Read carefully the given paragraph and answer the following questions.

- i. The product of the powers of each prime factors of 36 is:

- a. 2 b. 4 c. 3 d. 5

- ii. If p and q are positive integers such that $p = ab^2$ and $q = a^2b$, where a, b are prime numbers, then the LCM (p, q) is:
- a. ab b. a^2b^2 c. a^3b^2 d. a^3b^3
- iii. What is the minimum number of books you will acquire for the class library, so that they can be distributed equally among students of Section A or Section B?
- a. 144 b. 128 c. 288 d. 272
- iv. If the product of two positive integers is equal to the product of their HCF and LCM is true, then the HCF (32, 36) is:
- a. 2 b. 4 c. 6 d. 8
- v. If 288 books are distributed among the students of section B, how many will each get?
- a. 9 b. 8 c. 6 d. 7

CHAPTER - 2 (POLYNOMIALS)

Solve the following questions:

Q1. Graph of a quadratic polynomial is:

- a. straight line b. circle c. parabola d. ellipse

Q2. The parabola representing a quadratic polynomial $f(x) = ax^2 + bx + c$ opens upward when:

- a. $a > 0$ b. $a < 0$ c. $a = 0$ d. $a > 1$

Q3. The parabola representing a quadratic polynomial $f(x) = ax^2 + bx + c$ opens downward when:

- a. $a < 0$ b. $a > 0$ c. $a < 1$ d. $a > 1$

Q4. If one root of the polynomial $f(x) = 3x^2 + 11x + p$ is reciprocal of the other, then the value p is:

- a. 0 b. 3 c. $\frac{1}{3}$ d. -3

Q5. The zeroes of the quadratic polynomial $x^2 + kx + k$, where $k > 0$:

- a. are both positive b. are both negative
c. are always equal d. are always unequal

Q6. The graph of a quadratic polynomial intersects the X-axis at most at _____.

- a. 1 point b. 2 points c. 3 points d. 0 point

Q7. If the zeroes of the quadratic polynomial $x^2 + (a + 1)x + b$ are 2 and -3, then:

- a. $a = -7, b = -1$ b. $a = 5, b = -1$ c. $a = 2, b = -6$ d. $a = 0, b = -6$

Q8. The zeroes of the polynomials $x^2 + x - 2$ are:

- a. 1,2 b. -1,-2 c. 1,-2 d. -1,2

Q9. If 2 and $\frac{1}{2}$ are the zeroes of $px^2 + 5x + r$, then:

- a. $p = r = -2$ b. $p = r = 2$ c. $p = 2, r = -2$ d. $p = -2, r = 2$

Q10. If one of the zeroes of the quadratic polynomials $x^2 + 3x + k$ is 2, then the value of k is:

- a. 10 b. -10 c. -7 d. -2

- Q11.** How many maximum number of zeroes can a polynomial of degree 3 have?
- Q12.** What is the degree of the polynomial $(x+1)(x^2 - x - x^4 + 1)$?
- Q13.** The sum and product of zeroes of $p(x) = 63x^2 - 7x - 9$ are S and P respectively. Find $27S + 14P$.
- Q14.** Write a quadratic polynomial having zeroes 1 and -2.
- Q15.** If the sum of the zeroes of the quadratic polynomial $3x^2 - kx + 6$ is 3, then find the value of k.
- Q16.** Find the quadratic polynomial, whose zeroes are $5 - 3\sqrt{2}$ and $5 + 3\sqrt{2}$.
- Q17.** If one zero of the polynomial $x^2 + a$ is -3, then find its other zero.
- Q18.** Show that $\frac{1}{2}$ and $-\frac{3}{2}$ are the zeroes of the polynomial $4x^2 + 4x - 3$ and verify the relationship between zeroes and coefficients of the polynomial.
- Q19.** If the sum of the zeroes of the quadratic polynomial $(p^2 - 23)x^2 - 2x - 12$ is 1, then find the value of p.
- Q20.** Find the zeroes of the polynomials $7y^2 - \frac{11}{3}y - \frac{2}{3}$ and verify the relationship between the coefficients and the zeroes of the polynomials.
- Q21.** If α and β are the zeroes of a quadratic polynomial such that $\alpha + \beta = 24$ and $\alpha - \beta = 8$. Find the quadratic polynomial having α and β as its zeroes. Verify the relationship between the zeroes and coefficients of the polynomial.
- Q22.** If α and β are the zeroes of the quadratic polynomial $p(x) = x^2 + x - 2$, find the value of $\frac{1}{\alpha} - \frac{1}{\beta}$
- Q23.** If α and β are the zeroes of the quadratic polynomial $p(t) = t^2 - 4t + 3$, find the value of $\alpha^4\beta^3 + \alpha^3\beta^4$

Case-study based questions:

- Q24.** The below pictures show few natural examples of parabolic shape which can be represented by a quadratic polynomial. In structures, their curve represents an efficient method of load, and so can be found in bridges and in architecture in a variety of forms.



Based on the above information, answer the following questions:

- i. In the standard form of quadratic polynomial, $ax^2 + bx + c$, a, b and c:
 - a. all are real numbers
 - b. all are rational numbers
 - c. 'a' is a non-zero real number and b and c are any real numbers.
 - d. all are integers

- ii. The shape of the graph represented by quadratic polynomial is a:
- a. Parabola b. Straight line c. Circle d. Ellipse
- iii. If the polynomial is $x^2 - 3x - 10$, then its zeroes:
- a. 2, 5 b. 2, -5 c. -2, -5 d. none of these
- iv. The graph of $x^2 + 4 = 0$
- a. intersects X - axis at two distinct points
 b. touches X - axis at a point
 c. neither touches nor intersects X - axis
 d. either touches or intersects X - axis
- v. If the sum of the roots is $-p$ and product of the roots $-\frac{1}{p}$ is, then the quadratic polynomial is:
- a. $k\left(-px^2 + \frac{x}{p} + 1\right)$ b. $k\left(px^2 - \frac{x}{p} - 1\right)$
 c. $k\left(x^2 + px - \frac{1}{p}\right)$ d. $k\left(x^2 - px + \frac{1}{p}\right)$

CHAPTER - 3 (PAIR OF LINEAR EQUATIONS IN TWO VARIABLES)

Solve the following questions:

- Q1.** 5 chairs and 4 tables together cost Rs. 2800 while 4 chairs and 3 tables together cost Rs. 2170. Algebraic representation of the situation can be:
- a. $5x - 4y = 2800, 4x - 3y = 2170$ b. $5x + 4y = 2800, 4x + 3y = 2170$
 c. $5x + 3y = 2800, 4x + 3y = 2170$ d. $5x - 3y = 2800, 4x - 3y = 2170$
- Q2.** The sum of the numerator and denominator of a fraction is 8. If the denominator is increased by 1, the fraction becomes $\frac{1}{2}$. Algebraic representation of the situation can be:
- a. $x + y = 8, \frac{x+1}{y} = \frac{1}{2}$ b. $x + y = 8, \frac{x}{y} + 1 = \frac{1}{2}$
 c. $\frac{x}{y} = 8, \frac{x}{y+1} = \frac{1}{2}$ d. $x + y = 8, \frac{x}{y+1} = \frac{1}{2}$
- Q3.** The age of a daughter is one-third the age of her mother. If the present age of mother is x years, then the age (in years) of the daughter after 15 years is:
- a. $\frac{x}{3} + 15$ b. $\frac{x+15}{3}$ c. $x + 5$ d. $\frac{x}{3} - 15$
- Q4.** If $x = a, y = b$ is the solution of the pair of equation $x - y = 2$ and $x + y = 4$, then the respective values of a and b are:
- a. 3, 5 b. 5, 3 c. 3, 1 d. -1, -3

- iv. How many point(s) lie on the line $x - 3y = 2$?
- a. one b. two c. three d. infinitely
- v. If the line $2x + 6y = 5$ intersect the X - axis, then find its coordinate:
- a. $(-2.5, 0)$ b. $(2.5, 0)$ c. $(0, 2.5)$ d. $(0, -2.5)$

CHAPTER - 4 (QUADRATIC EQUATIONS)

Solve the following questions:

- Q1.** If $\frac{1}{2}$ is a root of the equation $x^2 + kx - \frac{5}{4} = 0$, then the value of k is:
- a. 2 b. -2 c. $\frac{1}{4}$ d. $\frac{1}{2}$
- Q2.** Which of the following equations has the sum of its roots as 3?
- a. $2x^2 - 3x + 6 = 0$ b. $-x^2 + 3x - 3 = 0$
- c. $\sqrt{2}x^2 - \frac{3}{\sqrt{2}}x + 1 = 0$ d. $3x^2 - 3x + 3 = 0$
- Q3.** Which of the following equations has 2 as a root?
- a. $x^2 - 4x + 5 = 0$ b. $-x^2 + 3x - 12 = 0$
- c. $2x^2 - 7x + 6 = 0$ d. $3x^2 - 6x - 2 = 0$
- Q4.** Value of k for which the quadratic equation $2x^2 - kx + k = 0$ has equal roots is:
- a. 0 only b. 4 c. 8 only d. 0, 8
- Q5.** $(x^2 + 1)^2 - x^2 = 0$ has _____.
- a. four real roots b. two real roots c. no real roots d. one real root
- Q6.** Which of the following equations has no real roots?
- a. $x^2 - 4x + 3\sqrt{2} = 0$ b. $x^2 + 4x - 3\sqrt{2} = 0$ c. $x^2 - 4x - 3\sqrt{2} = 0$ d. $3x^2 + 4\sqrt{3} + 4 = 0$
- Q7.** The quadratic equation $2x^2 - \sqrt{5}x + 1 = 0$:
- a. two distinct real roots b. two equal real roots
- c. no real roots d. more than two real roots
- Q8.** Is -8 is a solution of the equation $3x^2 + 8x + 2 = 0$?
- a. Yes b. No
- c. Cannot be determined d. None of these
- Q9.** If the roots of the quadratic equation $4x^2 + px + 9 = 0$ are equal, then the value of p is:
- a. ± 9 b. ± 6 c. ± 12 d. ± 3
- Q10.** If $x = 2$ is a root of both the equation $3x^2 + 2x + a = 0$ and $bx^2 + 2x + a = 0$, then ab is _____.
- a. 40 b. 32 c. 22 d. 15
- Q11.** Find the value of k for which -1 is a root of the quadratic equation $kx^2 + x - 6 = 0$.
- Q12.** Find the roots of the equation $x^2 + 7x + 10 = 0$.
- Q13.** For what values of k, the quadratic equation $kx(x-2) = 0$ has real roots?

Q14. If one root of the quadratic equation $5x^2 + 13x + k = 0$ is reciprocal of the other, then find the value of k .

Q15. Find the values of k for which the quadratic equation $9x^2 - 3kx + k = 0$ has real roots.

Q16. The sum of the squares of two consecutive multiples of 7 is 637. Find the multiples.

Q17. Solve the quadratic equation by factorisation method: $\frac{2x}{x-3} + \frac{1}{2x+3} + \frac{3x+9}{(x-3)(2x+3)} = 0$

Q18. Find the value of k for which $x^2 - 2x(1 + 3k) + 7(3 + 2k) = 0$ has equal roots.

Q19. Solve by factorization: $a^2 b^2 x^2 + b^2 x - a^2 x - 1 = 0$

Q20. The sum of two number is 11 and the sum of their reciprocals is $\frac{11}{28}$. Find the numbers.

Q21. A two-digit number is such that the product of the digits is 12. When 36 is added to the number the digits interchange their places. Find the two-digit number.

Q22. Find the roots of the equation $ax^2 + a = a^2x + x$.

Q23. Solve for x , $\sqrt{6}x + 7 - (2x - 7) = 0$

Case-study based questions:

Q24. Amit is preparing for his upcoming semester exam. For this, he has to practice the chapter of Quadratic Equations. So he started with factorization method. Let two linear factors of $ax^2 + bx + c$ be $(px + q)$ and $(rx + s)$.

Therefore, $ax^2 + bx + c = (px + q)(rx + s) = prx^2 + (ps + qr)x + qs$.

Now, factorize each of the following quadratic equations and find the roots.

i. $6x^2 + x - 2 = 0$

- a. 1, 6 b. $\frac{1}{2}, \frac{-2}{3}$ c. $\frac{1}{3}, \frac{-1}{2}$ d. $\frac{3}{2}, -2$

ii. $2x^2 + x - 300 = 0$

- a. 30, -60 b. $60, -\frac{2}{5}$ c. $12, \frac{-25}{2}$ d. none of these

iii. $x^2 - 8x + 6 = 0$

- a. 3, 3 b. 3, -3 c. 4, -4 d. 4, 4

iv. $6x^2 - 13x + 5 = 0$

- a. $2, \frac{3}{5}$ b. $\frac{-5}{3}, -2$ c. $\frac{1}{2}, \frac{-3}{5}$ d. $\frac{1}{2}, \frac{5}{3}$

v. $100x^2 - 20x + 1 = 0$

- a. $\frac{1}{10}, \frac{1}{10}$ b. -10, -10 c. -10, $\frac{1}{10}$ d. $\frac{-1}{10}, \frac{-1}{10}$

CHAPTER - 5 (ARITHMETIC PROGRESSIONS)

Solve the following questions:

Q1. In an AP, if $d = -4$, $n = 7$, $a_n = 4$, then a is:

- a. 6 b. 7 c. 20 d. 28


Q2. The first term of an AP is p and common difference is q , then its 10th term is:

- a. $q + 9p$ b. $p - 9q$ c. $p + 9q$ d. $2p + 9q$

- Q3.** The 21st of an AP whose first two terms are -3 and 4 is:
 a. 17 b. 137 c. 143 d. -143
- Q4.** If 7 times the 7th term of an AP is equal to 11 times its 11th term, then its 18th term will be:
 a. 7 b. 11 c. 18 d. 0
- Q5.** The sum of first 16 terms of the A.P. 10, 6, 2,..... is:
 a. -320 b. 320 c. -352 d. -400
- Q6.** In an AP, if $a = 1$, $a_n = 20$ and $S_n = 399$, then n is:
 a. 19 b. 21 c. 38 d. 42
- Q7.** The sum of first five multiple of 3 is:
 a. 45 b. 55 c. 65 d. 75
- Q8.** For the AP: $\frac{3}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{3}{2}, \dots$ write the first term and common difference.
- Q9.** Find the common difference of the AP:
 $\frac{1}{a}, \frac{3-a}{3a}, \frac{3-2a}{3a}, \dots, (a \neq 0)$.
- Q10.** Find the 12th term of the AP: 5, 8, 11, 14,
- Q11.** Find the sum of all even natural number less than 100.
- Q12.** Find the sum of first n terms of the series $\sqrt{2} + \sqrt{8} + \sqrt{18} + \dots$
- Q13.** Which term of the AP: 121, 117, 113, ... is its first negative term?
- Q14.** The 17th term of an AP is 5 more than twice its 8th term. If the 11th term of the AP is 43, then find its n^{th} term.
- Q15.** The ratio of the 5th and 3rd terms of an AP is 2 : 5. Find the ratio of the 15th and 7th terms.
- Q16.** The sum of the first three terms of an A.P. is 33. If the product of first and third term exceeds the second term by 29, find the AP.
- Q17.** An A.P. consists of 37 terms. The sum of the three middle most terms is 225 and the sum of the last three terms is 429. Find the AP.
- Q18.** A manufacturer of TV sets produced 600 units in the third year and 700 units in the seventh year. Assuming that the production increases uniformly by a fixed number every year, find the production in
 i. the first year ii. the 10th year iii. 7 years
- Q19.** The sum of four consecutive numbers in an AP is 32 and the ratio of the product of the first and the last term to the product of two middle terms is 7 : 15. Find the numbers.
- Q20.** The sum of first six terms of an AP is 42. The ratio of 10th term to its 30th term is 1 : 3. Calculate the first term and 13th term of an AP.

Case-study based questions:

- Q21.** Your friend Veer wants to participate in a 200 m race. He can currently run that distance in 51 sec. and with each day of practice it takes him 2 sec. less. He wants to do in 31 sec.




Based on the above information, answer the following questions:

- i. Which of the following terms are in AP for the given situation?
 - a. 51, 53, 55....
 - b. 51, 49, 47....
 - c. -51, -53, -55....
 - d. 51, 55, 59...
- ii. What is the minimum number of days he needs to practice till his goal is achieved?
 - a. 10
 - b. 12
 - c. 11
 - d. 9
- iii. Which of the following term is not in the AP of the above given situation?
 - a. 41
 - b. 30
 - c. 37
 - d. 39
- iv. If n^{th} term of an AP is given by $a_n = 2n + 3$ then common difference of an AP is:
 - a. 2
 - b. 3
 - c. 5
 - d. 1
- v. The value of x , for which $2x, x+10, 3x+2$ are three consecutive terms of an AP:
 - a. 6
 - b. -6
 - c. 18
 - d. -18

CHAPTER - 6 (TRIANGLES)

Solve the following questions:

- Q1.** The height of mountains is found out using the idea of indirect measurements which is based on the_____.
- a. principle of congruent figures
 - b. principle of similarity of figures
 - c. principle of equality of figures
 - d. None of the above
- Q2.** If triangle ABC is similar to triangle DEF such that $3AB = DE$ and $BC = 9\text{cm}$, then EF is equal to:
- a. 27 cm
 - b. 3 cm
 - c. 6 cm
 - d. 9 cm
- Q3.** In the given figure, value of x (in cm) is:




- a. 4
- b. 5
- c. 6
- d. 8

Q4. "If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio." This theorem is known as_____.

- a. Pythagoras Theorem
- b. Laplace Theorem
- c. Thales Theorem
- d. Area Theorem

Q5. In the given figure, if $AB \parallel DC$, find the value of x :



- a. 5
- b. 7
- c. 6
- d. 4

Q6. In triangles ABC and DEF, $\angle B = \angle E, \angle F = \angle C$ and $AB = 3 DE$. Then, the two triangles are:

- a. congruent but not similar
- b. similar but not congruent
- c. neither congruent nor similar
- d. congruent as well as similar

Q7. If perimeter of a triangle is 100 cm and the length of two sides are 30 cm and 40 cm, the length of third side will be:

- a. 30 cm
- b. 40 cm
- c. 50 cm
- d. 60 cm

Q8. If triangles ABC and DEF are similar and $AB = 4$ cm, $DE = 6$ cm, $EF = 9$ cm and $FD = 12$ cm, the perimeter of triangle is:

- a. 22 cm
- b. 20 cm
- c. 21 cm
- d. 18 cm


Q9. It is given that $\triangle ABC \sim \triangle DEF, \angle A = 30^\circ, \angle C = 50^\circ, AB = 5$ cm, $AC = 8$ cm and $DF = 7.5$ cm. Then, the following is true:

- a. $DE = 12$ cm, $\angle F = 50^\circ$
- b. $DE = 12$ cm, $\angle F = 100^\circ$
- c. $EF = 12$ cm, $\angle D = 100^\circ$
- d. $EF = 12$ cm, $\angle D = 30^\circ$

Q10. Areas of an equilateral triangle with side length 'a' is equal to:


- a. $\sqrt{3}/2a$
- b. $\sqrt{3}/2a^2$
- c. $\sqrt{3}/4a^2$
- d. $\sqrt{3}/4a$

Q11. If the given figure, $DE \parallel BC, AB = 5.6$ cm and $AD = 1.6$ cm, then find $AE : EC$.




Q12. If in $\triangle ABC, AB = 6$ cm and $DE \parallel BC$ such that $AE = \frac{1}{4} AC$, then find the length of AD.


- Q13.** In the $\triangle ABC$, D and E are points on side AB and AC respectively such that $DE \parallel BC$. If $AE = 2$ cm, $AD = 3$ cm and $BD = 4.5$ cm then find CE.
- Q14.** If one diagonal of a trapezium divides the other diagonal in the ratio 1:2, prove that one of the parallel sides is double the other.
- Q15.** In two triangles, it is given that the corresponding angles are equal. State whether the two triangles are congruent, similar or both.
- Q16.** If a line is drawn parallel to one side of a triangle, to intersect the other two sides in distinct points, the other two sides are divided in the same ratio.
- Q17.** In figure, $\triangle FEC \cong \triangle GBD$ and $\angle 1 = \angle 2$. Prove that $\triangle ADE \cong \triangle ABC$.




- Q18.** In the given figure, OB is the perpendicular bisector of the line segment DE, $FA \perp OB$ and FE intersects OB at the points C. Prove that $\frac{1}{OA} + \frac{1}{OB} = \frac{2}{OC}$.



- Q19.** A vertical pole which is 2.25 m long casts a 6.75 m long shadow on the ground. At the same time, a vertical tower casts a 90 m long shadow on the ground. Find the height of the tower.
- Q20.** In the given figure, find the value of x in terms of a, b and c.




Q21. In the given figure, $DB \perp BC$, $AC \perp BC$ and $DE \perp AB$. Prove that $\frac{BE}{DE} = \frac{AC}{BC}$.



Case-study based questions:

Q22. Vijay is trying to find the average height of a tower near his house. He is using the properties of similar triangles. The height of Vijay's house is 20 m when Vijay's house casts a shadow 10 m long on the ground. At the same time, the tower casts a shadow 50 m long on the ground and the house of Ajay casts 20 m shadow on the ground.



Based on the above information, give the answer of the following questions:

- i. What is the height of the tower?
 - a. 20 m
 - b. 50 m
 - c. 100 m
 - d. 200 m
- ii. What will be the length of the shadow of the tower when Vijay's house casts a shadow of 12 m?
 - a. 75 m
 - b. 50 m
 - c. 45 m
 - d. 60 m
- iii. What is the height of Ajay's house?
 - a. 30 m
 - b. 40 m
 - c. 50 m
 - d. 20 m
- iv. When the tower casts a shadow of 40 m, same time what will be the length of the shadow of Ajay's house?
 - a. 16 m
 - b. 32 m
 - c. 20 m
 - d. 8 m
- v. When the tower casts a shadow of 40 m, same time what will be the length of the shadow of Vijay's house?
 - a. 15 m
 - b. 32 m
 - c. 16 m
 - d. 8 m

CHAPTER - 7 (COORDINATE GEOMETRY)

Solve the following questions:


- Q1.** Distance of the point (5, -4) from X - axis is _____.
- a. 5 units b. 4 units c. 1 unit d. 9 units
- Q2.** If the point (x, y) is equidistant from the point (2, 1) and (1, -2), then:
- a. $x + 3y = 0$ b. $3x + y = 0$ c. $x + 2y = 0$ d. $3x + 2y = 0$
- Q3.** If A(5, 3), B(11, -5) and P(12, y) are the vertices of a right triangle, right angled at P, then y:
- a. -2 or 4 b. -2 or -4 c. 2 or -4 d. 2 or 4
- Q4.** The x-coordinate of a point P is twice its y-coordinate. If P is equidistant from Q(2, -5) and R(-3, 6), then the coordinates of P are:
- a. (16, 8) b. (14, 7) c. (18, 9) d. (10, 5)
- Q5.** The equation of the perpendicular bisector of line segment joining points A(4, 5) and B(-2, 3) is:
- a. $2x - y + 7 = 0$ b. $3x + 2y - 7 = 0$ c. $3x - y - 7 = 0$ d. $3x + y - 7 = 0$
- Q6.** The point which divides the line segment joining the points (7, -6) and (3, 4) in ratio 1:2 internally lies in the _____.
- a. I quadrant b. II quadrant c. III quadrant d. IV quadrant
- Q7.** The ratio in which the line segment joining (1, -5) and (-4, 5) is divided by the x - axis is:
- a. 1 : 1 b. 1 : 2 c. 2 : 1 d. None of these
- Q8.** The mid - point of the line segment AB is P(0, 4). If the coordinates of B are (-2, 3) then the coordinates of A are:
- a. (2, 5) b. (-2, -5) c. (2, 9) d. (-2, 11)
- Q9.** Find the point on x-axis which is equidistant from the points (2, -2) and (-4, 2).
- Q10.** Find a point on the y-axis which is equidistant from (6, 5) and (-4, 3).
- Q11.** If point P divides the line segment joining the points A(2, 1) and B(6, 5) in the ratio 1:3, then find the coordinates of P.
- Q12.** If the center of a circle is $\left(\frac{3}{2}, \frac{-5}{2}\right)$ and one end of the diameter is (2, 1), then find the coordinates of the other end.
- Q13.** If the mid-point of the line segment joining the points P(6, b-2) and Q(-2, 4) is (2, -3), find the value of b.
- Q14.** Find the points on the x-axis which is equidistant from (2, -5) and (-2, 9).
- Q15.** Find the ratio in which the line $2x + y = 4$ divides the join of A(2, -2) and B(3, 7). Also, find the coordinates of the point of their intersection.
- Q16.** The three vertices of a parallelogram are (6, 1), (8, 2) and (9, 4). Find the fourth vertex.
- Q17.** The line segment joining the points (3, -4) and (1, 2) is trisected at the points P and Q. If the coordinates of P and Q are (a, -2) and $\left(\frac{5}{3}, b\right)$ respectively, find the values of a and b.
- Q18.** Find the coordinates of the points which divide the line segment joining A(-2, 2) and B(2, 8) into four equal parts.

Q19. Find the ratio in which the line $2x + 3y - 5 = 0$ divides the line segment joining the points $(8, -9)$ and $(2, 1)$. Also find the coordinates of the point of division.

Q20. Show that the points $(1, 7)$, $(4, 2)$, $(-1, -1)$ and $(-4, 4)$ are the vertices of a square.

Case-study based questions:

Q21. Sonu went to the lab near to his home for COVID 19 test along with his family members. The seats in the waiting area were as per the norms of distancing during this pandemic (as shown in the above figure). His family members took their seats surrounded by black circular area.



Based on the above information give the answer of the following questions.

- i. Considering O as the origin, what are the coordinates of seat of Sonu and Goru respectively?
 - a. $(2, 3)$ and $(6, 5)$
 - b. $(3, 2)$ and $(5, 6)$
 - c. $(3, 6)$ and $(5, 2)$
 - d. $(6, 3)$ and $(2, 5)$
- ii. What is the distance between Golu and Monu?
 - a. $\sqrt{5}$ units
 - b. $2\sqrt{5}$ units
 - c. $3\sqrt{5}$ units
 - d. $4\sqrt{5}$ units
- iii. What will be the coordinates of a point exactly between Sonu and Goru where a person can be seated?
 - a. $\left(\frac{5}{2}, \frac{9}{2}\right)$
 - b. $\left(\frac{11}{2}, \frac{7}{2}\right)$
 - c. $(4, 4)$
 - d. $\left(\frac{9}{2}, 4\right)$
- iv. Find the area covered by Sonu and its members, if all four seats connected with a rope.
 - a. $2\sqrt{5}$ sq. units
 - b. $\sqrt{10}$ sq. units
 - c. $2\sqrt{10}$ sq. units
 - d. 10 sq. units
- v. If the doctor divides the rope joining Sonu and Goru in the ratio $1 : 2$, then the coordinates of the seat of the doctor is:
 - a. $\left(\frac{10}{3}, \frac{11}{3}\right)$
 - b. $\left(\frac{11}{3}, \frac{10}{3}\right)$
 - c. $(4, 4)$
 - d. $(3, 4)$

CHAPTER - 8 (INTRODUCTION TO TRIGONOMETRY)

Q1. The maximum value of $\sin \theta$ is _____.

- a. $\frac{1}{2}$
- b. $\frac{\sqrt{3}}{2}$
- c. 1
- d. $\frac{1}{\sqrt{2}}$

Q2. If $\tan \theta = \frac{3}{4}$, then $\cos^2 \theta - \sin^2 \theta =$

- a. $\frac{7}{25}$ b. 1 c. $\frac{-7}{25}$ d. $\frac{4}{25}$

Q3. If $\tan \theta = \frac{a}{b}$, then $\frac{(a \sin \theta - b \cos \theta)}{(a \sin \theta + b \cos \theta)} =$

- a. $\frac{(a^2 + b^2)}{(a^2 - b^2)}$ b. $\frac{(a^2 - b^2)}{(a^2 + b^2)}$ c. $\frac{a^2}{(a^2 + b^2)}$ d. $\frac{b^2}{(a^2 + b^2)}$

Q4. Match the columns:

| Column I | | Column II | |
|----------|---|-----------|---------------|
| 1. | Side opposite to angle θ ----- Hypotenuse | A. | $\tan \theta$ |
| 2. | Side adjacent to angle θ ----- Hypotenuse | B. | $\sin \theta$ |
| 3. | Side opposite to angle θ ----- Side adjacent to angle θ | C. | $\cos \theta$ |

- a. 1-A, 2-C, 3-B b. 1-B, 2-C, 3-A c. 1-A, 2-B, 3-C d. 1-C, 2-B, 3-A

Q5. $\sqrt{-4 + \sqrt{8 + 16 \operatorname{cosec}^4 \theta + \sin^4 \theta}} = A \operatorname{cosec} \theta + B \sin \theta$, then A and B are:

- a. 2 and -1 b. 1 and $\frac{1}{2}$ c. $\frac{1}{2}$ and $\frac{1}{3}$ d. 3 and -4

Q6. Value of $\frac{\sin 60^\circ + \cot 45^\circ - \operatorname{cosec} 30^\circ}{\sec 60^\circ - \cos 30^\circ + \tan 45^\circ}$ is _____.

- a. $\frac{4\sqrt{3} - 9}{33}$ b. $\frac{4\sqrt{3} + 9}{33}$ c. $\frac{9\sqrt{3} - 4}{33}$ d. $\frac{9\sqrt{3} + 4}{33}$

Q7. If $\frac{x \operatorname{cosec}^2 30^\circ \sec^2 45^\circ}{8 \cos^2 45^\circ \sin^2 60^\circ} = \tan^2 60^\circ - \tan^2 30^\circ$, then x =

- a. 1 b. -1 c. 2 d. 0

Q8. If $\sin \alpha = \frac{1}{2}$ and $\tan \beta = \frac{1}{\sqrt{3}}$, $\alpha > 0^\circ$, $\beta > 0^\circ$ then the value of $\cot(\alpha + \beta)$ is:

- a. $\frac{1}{\sqrt{3}}$ b. $\sqrt{3}$ c. 1 d. 0

Q9. If $a \cos \theta + b \sin \theta = m$ and $a \sin \theta - b \cos \theta = n$, then $a^2 + b^2 =$

- a. $m^2 - n^2$ b. $n^2 - m^2$ c. $m^2 + n^2$ d. $m^2 n^2$

Q10. If $\tan \theta + \sin \theta = m$ and $\tan \theta - \sin \theta = n$, then $m^2 - n^2$ is equal to:

- a. \sqrt{mn} b. $\sqrt{\frac{m}{n}}$ c. $4\sqrt{mn}$ d. None of these

Q11. If $3 \cot \theta = 2$, then what is the value of $\tan \theta$?

Q12. If $\cot A = \frac{12}{5}$, find the value of $(\sin A + \cos A) \operatorname{cosec} A$.

Q13. Find the value of $\cos 60^\circ \cos 30^\circ + \sin 60^\circ \sin 30^\circ$.

Q14. If $5 \tan \theta = 4$, then find the value of $\frac{5 \sin \theta - 3 \cos \theta}{5 \sin \theta + 3 \cos \theta}$

Q15. Verify: $4(\sin^2 30^\circ + \cos^2 60^\circ) - 3(\cos^2 45^\circ - \sin^2 90^\circ) = 2$

Q16. If $\sin (A - B) = \frac{1}{2}$, $\cos (A + B) = \frac{1}{2}$, find A and B.

Q17. Prove that: $(1 + \cot A - \operatorname{cosec} A)(1 + \tan A + \sec A) = 2$

Q18. Evaluate: $\frac{\tan^2 60^\circ + 4 \sin^2 45^\circ + 3 \sec^2 30^\circ + 5 \cos^2 90^\circ}{\operatorname{cosec} 30^\circ + \sec 60^\circ - \cot^2 30^\circ}$

Q19. Prove that: $\frac{1}{\operatorname{cosec} \theta - \cot \theta} - \frac{1}{\sin \theta} = \frac{1}{\sin \theta} - \frac{1}{\operatorname{cosec} \theta + \cot \theta}$


Q20. Prove that: $\sec^4 A - \sec^2 A = \tan^4 A + \tan^2 A$

Q21. Prove that: $\frac{\sin \theta}{\cot \theta + \operatorname{cosec} \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta}$

Q22. Prove that: $\frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} = 1 + \sec \theta \cdot \operatorname{cosec} \theta$

Case-study based questions:

Q23. A sailing boat with triangular masts is shown below. Two right triangles can be observed. Triangles PQR and PQS, both right - angled at Q. The distance QR = 2m and QS = 3m and height PQ = 5m.



Based on the above information, give the answer of the following questions :

i. The value of $\sec S$ is:

- a. $\frac{\sqrt{34}}{5}$ b. $\frac{\sqrt{34}}{3}$ c. $\frac{5}{3}$ d. $\frac{3}{\sqrt{34}}$

ii. The value of $\operatorname{cosec} R$ is:

- a. $\frac{\sqrt{29}}{5}$ b. $\frac{\sqrt{29}}{2}$ c. $\frac{2}{5}$ d. $\frac{5}{\sqrt{29}}$

iii. The value of $\tan S + \cot R$ is:

- a. $\frac{9}{4}$ b. $\frac{5}{3}$ c. $\frac{2}{5}$ d. $\frac{31}{15}$

iv. Value of $\sin^2 R - \cos^2 S$ is:

- a. 0 b. 1 c. $\frac{97}{85}$ d. $\frac{589}{986}$

v. Value of $\sin^2 S - \cos^2 R$ is:

- a. 0 b. 1 c. $\frac{97}{85}$ d. $\frac{589}{986}$

CHAPTER - 9 (SOME APPLICATIONS OF TRIGONOMETRY)

Solve the following questions:

Q1. A ladder of 10 m length touches a wall at height of 5 m. The angle θ made by it with the horizontal is:

- a. 60° b. 90° c. 30° d. 45°

Q2. The measure of angle of elevation of top of tower $75\sqrt{3}$ m high from a point at a distance of 75 m from foot of tower in a horizontal plane is:

- a. 30° b. 60° c. 90° d. 45°

Q3. A pole 10 m high cast a shadow 10 m long on the ground, then the sun's elevation is_____.

- a. 60° b. 45° c. 30° d. 90°

Q4. If the ratio of height of a tower and the length of its shadow on the ground is $\sqrt{3} : 1$, then the angle of elevation of the sun is:

- a. 60° b. 45° c. 30° d. 90°

Q5. If the angle of depression of an object from a 75 m high tower is 30° , then the distance of the object from base of tower is_____.

- a. 60° b. 45° c. 30° d. 90°

Q6. When the angle of elevation of sun is 30° the length of the shadow cast by 50 m high building is:

- a. $\frac{50}{\sqrt{3}}$ m b. $50\sqrt{3}$ m c. $25\sqrt{3}$ m d. $100\sqrt{3}$ m

Q7. If $AB = 4$ m and $AC = 8$ m, then angle of elevation of A as observed from C is:

- a. 60° b. 45° c. 30° d. 90°

- Q8.** If the altitude of the sun is 60° , the height of a tower which casts a shadow of length 30 m is:
- a. $30\sqrt{3}$ m b. $\frac{30}{3}\sqrt{3}$ m c. $15\sqrt{3}$ m d. 15 m
- Q9.** The angle of elevation of the top of a tower from a point on the ground is 45° . If the observer is 42 m away from the foot of the tower, the height of the tower is:
- a. 63 m b. 21 m c. 84 m d. 42 m
- Q10.** A ladder makes an angle of 60° with the ground when placed against a wall. If the foot of the ladder is 2 m away from the wall, then find the length of the ladder.
- Q11.** The tops of two towers of height x and y , standing on level ground, subtend angles of 30° and 60° respectively at the centre of the line joining their feet, then find $x : y$.
- Q12.** If the height and the length of the shadow of a man are the same, then find the angle of elevation of the Sun.
- Q13.** An observer 1.5 m tall is 20.5 m away from a tower of 22 m high. Determine the angle of elevation of the tower from the eye of the observer.
- Q14.** A man on the top of the vertical tower observes a car moving at a uniform speed toward him. If it takes 12 min for the angle of depression to change from 30° to 45° , how soon after this, the car will reach the tower?
- Q15.** The angle of elevation of an aeroplane from point A on the ground is 60° . After flight of 15 seconds, the angle of elevation change to 30° . If the aeroplane is flying at a constant height of $500\sqrt{3}$ m, find the speed of the plane in km/hr.
- Q16.** The angle of elevation of a jet plane from a point A on the ground is 60° . After a flight of 30 seconds, the angle of elevation changes to 30° . If the jet plane is flying at a constant height of $3600\sqrt{3}$ m, find the speed of the jet plane.
- Q17.** A pole 5 m high is fixed at the top of a tower. The angle of depression of the top of the pole observed from a point A on the ground is 60° and the angle of depression of the point A from the top of the tower is 45° . Find the height of the tower.
- Q18.** The angle of depression of the top and bottom of a building 50 metres high as observed from the top of a tower are 30° and 60° , respectively. Find the height of the tower and also the horizontal distance between the building and the tower.
- Q19.** A moving boat is observed from the top of a 150 m high cliff moving away from the cliff. The angle of depression of the boat changes from 60° to 45° in 2 minutes. Find the speed of the boat in m/h.
- Q20.** From a point in the ground, the angles of elevation of the bottom and top of a tower fixed at the top of a 20 m high building are 45° and 60° respectively. Find the height of the tower.
- Q21.** As observed from the top of a 100 m high lighthouse from the sea-level, the angles of depression of two ships are 30° and 45° . If one ship is exactly behind the other on the same side of the lighthouse, find the distance between the two ships. (Use $\sqrt{3} = 1.732$)

Case-study based questions:

- Q22.** A group of students of class X visited India Gate on an education trip. The teacher and students had interest in history as well. The teacher narrated that **India Gate**, official name **Delhi Memorial**, originally called **All-India War Memorial**, monumental sand stone arch in **New Delhi**, to the troops of British India who died in wars fought between 1914 and 1919. The teacher also said that India Gate, which is located at the eastern end of the Rajpath (formerly called the Kingsway), is about 138 feet (42 metres) in height.



Based on the above information, answer the following questions:

- i. What is the angle of elevation if they are standing at a distance of 42 m away from the monument?
 - a. 30°
 - b. 45°
 - c. 60°
 - d. 0°
- ii. They want to see the tower at an angle of 60° . So, they want to know the distance where they should stand and hence find the distance:
 - a. 25.24 m
 - b. 20.12 m
 - c. 42 m
 - d. 24.64 m
- iii. If the altitude of the Sun is at 60° , then the height of the vertical tower that will cast a shadow of length 20 m is:
 - a. $20\sqrt{3}$ m
 - b. $\frac{20}{\sqrt{3}}$ m
 - c. $\frac{15}{\sqrt{3}}$ m
 - d. $15\sqrt{3}$ m
- iv. The ratio of the length of a rod and its shadow is 1:1. The angle of elevation of the sun is:
 - a. 30°
 - b. 45°
 - c. 60°
 - d. 90°
- v. The angle formed by the line of sight with the horizontal when the object viewed is below the horizontal level is:
 - a. corresponding angle
 - b. angle of elevation
 - c. angle of depression
 - d. complete angle

CHAPTER-10 (CIRCLES)


Solve the following questions:

- Q1.** If two tangents inclined at an angle 60° are drawn to a circle of radius 3 cm, then length of each tangent is equal to:
 - a. $\frac{3}{2}\sqrt{3}$ cm
 - b. 6 cm
 - c. 3 cm
 - d. $3\sqrt{3}$ cm
- Q2.** The number of tangents that can pass through any point lying on a circle is_____.
 - a. 2
 - b. infinitely many
 - c. 1
 - d. none of these
- Q3.** From a point P which is at a distance of 13 cm from the centre O of a circle of radius 5 cm, the pair of tangents PQ and PR to the circle is drawn. Then, the area of the quadrilateral PQOR is:
 - a. 60 cm^2
 - b. 65 cm^2
 - c. 30 cm^2
 - d. 32.5 cm^2
- Q4.** Distance between two parallel lines is 14 cm. The radius of circle which will touch both of these lines is:
 - a. 6 cm
 - b. 7 cm
 - c. 12 cm
 - d. 14 cm

- Q5.** Angle between the tangent and the radius at the point of contact is_____.
- a. 0° b. 30° c. 60° d. 90°
- Q6.** If angle between two tangents drawn from a point P to a circle of radius a and centre O is 90° , then OP is:
- a. $\frac{\sqrt{3}a}{2}$ b. a c. $\sqrt{2}a$ d. $2a$
- Q7.** A tangent PQ at a point P of a circle of radius 6 cm meets a line through the centre O at a point Q, so that $OQ = 14$ cm, then length of PQ is:
- a. $4\sqrt{10}$ cm b. $6\sqrt{10}$ cm c. $5\sqrt{10}$ cm d. $7\sqrt{10}$ cm
- Q8.** How many tangents can a circle have from a point lying inside the circle?
- a. 2 b. infinitely many c. 1 d. none of these
- Q9.** From an external point P, tangents PA and PB are drawn to a circle with centre O. If $\angle PAB = 50^\circ$, then $\angle AOB$.
- Q10.** Find the length of the tangent drawn from a point P, 12 cm away from the center of a circle of radius 5 cm.
- Q11.** The length of the tangent from a point A to a circle of radius 9 cm is 12 cm. Find the distance of A from the center of the circle.
- Q12.** If two tangents inclined at an angle of 60° are drawn to a circle of radius 5 cm, then find the length of each tangent.
- Q13.** Prove that the tangents drawn from an external point to a circle are equal in length.
- Q14.** From a point Q, 13 cm away from the center of a circle, the length of tangent PQ to the circle is 12 cm. Find the radius of the circle.
- Q15.** PQ is a chord of length 8 cm of a circle of radius 5 cm. The tangents at P and Q intersect at T. Find the length of TP.
- Q16.** PA and PB are the tangents to a circle with center O. If $\angle APB = 70^\circ$. Find $\angle POA$.
- Q17.** Find the distance between two parallel tangents to a circle of radius 5 cm.
- Q18.** The length of tangent from a point A at distance 5 cm from the center of the circle is 4 cm. Find radius.
- Q19.** A circle is touching the side BC of $\triangle ABC$ at P and touching AB and AC produced at Q and R respectively. Prove that $AQ = \frac{1}{2}(\text{Perimeter of } \triangle ABC)$.
- Q20.** Prove the tangent at any point of a circle is perpendicular to the radius through the point of contact.

Case-study based questions:

- Q21.** A Ferris wheel (or a big wheel in the United Kingdom) is an amusement ride consisting of a rotating upright wheel with multiple passenger-carrying components (commonly referred to as passenger cars, cabins, tubs, capsules, gondolas, or pods) attached to the rim in such a way that as the wheel turns, they are kept upright, usually by gravity. After taking a ride in Ferris wheel, Aarti came out from the crowd and was observing her friends who were enjoying the ride . She was curious about the different angles and measures that the wheel will form. She forms the figure as given below.



Based on the above information, answer the following questions.

- i. In the given figure, find $\angle ROQ$:
 - a. 100
 - b. 150
 - c. 90
 - d. 61
- ii. Find $\angle RQP$:
 - a. 75
 - b. 60
 - c. 30
 - d. 90
- iii. Find $\angle RSQ$:
 - a. 60
 - b. 75
 - c. 100
 - d. 30
- iv. Find $\angle ORP$:
 - a. 90
 - b. 70
 - c. 100
 - d. 60

CHAPTER - 12 (AREAS RELATED TO CIRCLES)

Solve the following questions:

Q1. If the sum of the areas of two circles with radii R_1 and R_2 is equal to the area of a circle of radius R , then:

- a. $R_1 + R_2 = R$
- b. $R_1^2 + R_2^2 = R^2$
- c. $R_1 + R_2 < R$
- d. $R_1^2 + R_2^2 < R^2$

Q2. Match the columns

| | | | |
|----|------------------------------|----|---|
| 1. | Area of quadrant | A. | $\frac{1}{2} \pi r^2$ |
| 2. | Area of equilateral triangle | B. | $\frac{\sqrt{3}}{4} \times \text{side}^2$ |
| 3. | Area of semicircle | C. | $\pi r + 2r$ |
| 4. | Perimeter of semicircle | D. | $\frac{1}{4} \pi r^2$ |

- a. 1-A, 2-C, 3-D, 4-B
- b. 1-B, 2-C, 3-D, 4-B
- c. 1-D, 2-B, 3-A, 4-C
- d. 1-C, 2-B, 3-A, 4-D

Q3. The length of the minute hand of a clock is $\sqrt{21}$ cm. Find the area swept by the minute hand from 9 a.m. to 9.10 a.m.:

- a. 22 cm
- b. 11 cm²
- c. 45 cm²
- d. 31 cm²

Q4. A chord of a circle subtends an angle of 60° at the centre of the circle. If the length of the chord is 10 cm. Then the area of the major segment is: (Take $\pi = 3.14$ and $\sqrt{3} = 1.732$)

- a. 304.97 cm²
- b. 295 cm²
- c. 310 cm²
- d. 335 cm²


Q5. The area of the square that can be inscribed in a circle of radius 8 cm is_____.

- a. 256 cm^2 b. 128 cm^2 c. $64\sqrt{2} \text{ cm}^2$ d. 64 cm^2

Q6. The perimeter (in cm) of a square circumscribing a circle of radius a cm, is_____.


- a. $4a$ b. $5a$ c. $8a$ d. $10a$

Q7. In figure, two concentric circles with centre O, have radii 21cm and 42cm. If $\angle AOB = 60^\circ$, the area of the shaded region is:



- a. 3465 cm^2 b. 1295 cm^2 c. 2565 cm^2 d. 3980 cm^2

Q8. In the following figure, O is the centre of the circle with $AC = 24 \text{ cm}$, $AB = 7 \text{ cm}$ and $\angle BOD = 90^\circ$. The area of the shaded region is: (Use $\pi = 3.14$)



- a. 248 cm^2 b. 284 cm^2 c. 298 cm^2 d. 318 cm^2

Q9. Determine the length of an arc of quadrant of a circle of radius r.

Q10. In a circle of diameter 42 cm, if an arc subtends an angle of 60° at the center, find the length of an arc.

Q11. Find the area of the largest triangle that can be inscribed in a semicircle of radius r.


Q12. Find the area of the minor segment of a circle of radius 14 cm, when its central angle is 60° .

Q13. In a circle of radius 21 cm, an arc subtends an angle of 60° at the center. Find


- i. the area of the minor sector
- ii. length of an arc
- iii. perimeter of the sector.

Q14. Find the area of the minor segment of a circle of radius 42 cm, if length of the corresponding arc is 44 cm.


Q15. In the given figure, diameter of the largest semicircle is 108 cm and the diameter of the small circle is 36 cm. Find the area of the shaded region.




- Q16.** In the given figure, PQRS is a diameter of a circle of radius 6 cm. The lengths PQ, QR, and RS are equal. Semicircles are drawn on PQ and QS as diameters. Find the perimeter and area of the shaded region.




- Q17.** In figure, ABCD is a square of side 14 cm. Semi-circles are drawn with each side of square as diameter. Find the area of the shaded region. (use $\pi = \frac{22}{7}$)



- Q18.** Find the area of the shaded region in given figure, where arcs drawn with centres A, B, C and D intersect in pairs at mid-point P, Q, R and S of the sides AB, BC, CD and DA respectively of a square ABCD of side 12 cm.




- Q19.** In the given figure, shows a kite in which BCD is the shape of a quadrant of a circle of radius 42 cm. ABCD is a square and $\triangle CEF$ is an isosceles right angled triangle whose equal sides are 6 cm long. Find the area of the shaded region. (use $\pi = \frac{22}{7}$)



Case-study based questions:

Q20. We all love to eat pizzas, especially kids and a variety of pizzas are available in India which have been modified according to Indian taste and menu. From the Greeks to the Egyptians, from the Persians to the Indians, there have been incarnations of pizza served throughout history, Flatbreads, naan, and plakountos are all early preparations that could be considered cousins to the modern pizza, but there isn't a consensus as to which is first and whether these could even be considered precursors to pizza at all. Consider two pizzas, both of equal diameter, namely, 12 inches. The first pizza marked (I) has been cut into equal slices, whereas the second pizza, marked (II) has been cut into eight equal slices.



(I)

(II)

Based on the above information, answer the following questions.

- i. The area of one slice in pizza, marked (I), is:
 - a. 6π sq. inches
 - b. 8π sq. inches
 - c. 10π sq. inches
 - d. None of these
- ii. The perimeter of the pizza slice shown in (I), is:
 - a. $(\pi + 12)$ inch
 - b. $(\pi + 10)$ inch
 - c. $(2\pi + 10)$ inch
 - d. $(2\pi + 12)$ inch
- iii. The ratio of area of slice to the area of remaining pizza in (I), is:
 - a. 5 : 1
 - b. 1 : 5
 - c. 2 : 5
 - d. 5 : 3
- iv. The ratio of areas of each slice of pizza (I) and (II), is:
 - a. 3 : 4
 - b. 5 : 3
 - c. 4 : 3
 - d. 2 : 5
- v. The relation between area of a sector A, length of the arc l, angle θ subtended by the arc at the centre and radius of circle, is:
 - a. $\frac{1}{2}lr$
 - b. lr
 - c. $\frac{1}{3}lr$
 - d. $\frac{1}{2}lr^2$

CHAPTER- 13 (SURFACE AREAS AND VOLUMES)


Solve the following questions:

- Q1.** A cylindrical pencil sharpened at one edge is the combination of_____.
- a. a cone and a cylinder
 - b. frustum of a cone and a cylinder
 - c. a hemisphere and a cylinder
 - d. two cylinders
- Q2.** The length of the diagonal of a cube is $6\sqrt{3}$ cm. Its total surface area is:
- a. 144 cm^2
 - b. 216 cm^2
 - c. 180 cm^2
 - d. 108 cm^2

- Q3.** The curved surface area of a cylinder is 264 m^2 and its volume is 924 m^3 . The ratio of its diameter to its height is:
- a. 3 : 7 b. 7 : 3 c. 6 : 7 d. 7 : 6
- Q4.** A circular tent is cylindrical to a height of 4 m and conical above it. If its diameter is 105 m and its slant height is 40 m, then the total area of the canvas required is:
- a. 1760 m^2 b. 2640 m^2 c. 3960 m^2 d. 7920 m^2
- Q5.** A cubical ice-cream brick of edge 22 cm is to be distributed among some children by filling ice-cream cones of radius 2 cm and height 7 cm upto its brim. How many children will get ice-cream cones?
- a. 163 b. 263 c. 363 d. 463
- Q6.** If two solid hemispheres of same base radius r are joined together along their bases, then curved surface area of this new solid is:
- a. $4\pi r^2$ b. $6\pi r^2$ c. $3\pi r^2$ d. $8\pi r^2$
- Q7.** A medicine-capsule is in the shape of a cylinder of diameter 0.5 cm with two hemispheres stuck to each of its ends. The length of entire capsule is 2 cm. The capacity of the capsule is:
- a. 0.36 cm^3 b. 0.35 cm^3 c. 0.34 cm^3 d. 0.33 cm^3
- Q8.** A tank is made of the shape of a cylinder with a hemispherical depression at one end. The height of the cylinder is 1.45 m and radius is 30 cm. The total surface area of the tank is:
- a. 30 m b. 3.3 m c. 30.3 m d. 3300 m
- Q9.** A toy is in the form of a right circular with a hemisphere on the top. The radii of cylindrical and hemispherical parts are 5 cm each and height of cylindrical part is 14 cm. Then its curved surface area is:
- a. $190\pi \text{ cm}^2$ b. $140\pi \text{ cm}^2$ c. $175\pi \text{ cm}^2$ d. $185\pi \text{ cm}^2$
- Q10.** Find the slant height of a cone whose height is 4 cm and radius 3 cm.
- Q11.** Gayatri was making a mathematical model, in which she placed 4 cubes each of edge 20 cm one above another. Find the surface area of the resulting cuboid.
- Q12.** Water is flowing at the rate of 7 m/s through a circular pipe whose internal diameter is 2 cm into a cylindrical tank the radius of whose base is 40 cm. Determine the increase in the water level in $\frac{1}{2}$ hour.
- Q13.** If h , C and V respectively represent the height, curved surface area and volume of a cone, prove that $C^2 = \frac{3\pi Vh^3 + 9V^2}{h^2}$.
- Q14.** Two cones with same base radius 8 cm and height 15 cm are joined together along their bases. Find the surface area of the shape so formed.
- Q15.** A cylindrical tub, whose diameter is 12 cm and height 15 cm is full of ice-cream. The whole ice-cream is to be divided into 10 children in equal ice-cream cones, with conical base surmounted by hemispherical top. If the height of conical portion is twice the diameter of base, find the diameter of conical part of ice-cream cone.
- Q16.** A hemispherical depression is cut out from one face of a cubical wooden block of edge 21 cm, such that the diameter of the hemisphere is equal to edge of the cube. Determine the volume of the remaining block.

Case-study based questions:

Q17. On a Sunday, your Parents took you to a fair. You could see lot of toys displayed, and you wanted them to buy a RUBIK's cube and strawberry ice-cream for you.



Based on the above information, answer the following questions.

- i. The length of the diagonal if each edge measures 6cm is:
a. $3\sqrt{3}$ b. $3\sqrt{6}$ c. $\sqrt{12}$ d. $6\sqrt{3}$
- ii. Volume of the solid figure if the length of the edge is 7cm is:
a. 256 cm^3 b. 196 cm^3 c. 343 cm^3 d. 434 cm^3
- iii. What is the curved surface area of hemisphere (ice cream) if the base radius is 7cm?
a. 309 cm^2 b. 308 cm^2 c. 803 cm^2 d. 903 cm^2
- iv. Slant height of a cone, if the radius is 7cm and the height is 24 cm:
a. 26 cm b. 25 cm c. 52 cm d. 62 cm
- v. The total surface area of cone with hemispherical ice cream is:
a. 858 cm^2 b. 885 cm^2 c. 588 cm^2 d. 855 cm^2

CHAPTER - 14 (STATISTICS)

Solve the following questions:

- Q1.** Construction of a cumulative frequency table is useful in determining the:
- a. mean
 - b. median
 - c. mode
 - d. all the above three measures
- Q2.** The method used to find the mean of a given data is(are):
- a. direct method
 - b. assumed mean method
 - c. step deviation method
 - d. all the above
- Q3.** While computing mean of grouped data, we assume that the frequencies are:
- a. centred at the class marks of the classes
 - b. evenly distributed over all the classes
 - c. centred at the upper limits of the classes
 - d. centred at the lower limits of the classes
- Q4.** The empirical relationship between the three measures of central tendency is:
- a. $3 \text{ Median} = \text{Mode} + 2 \text{ Mean}$
 - b. $2 \text{ Median} = \text{Mode} + 2 \text{ Mean}$
 - c. $3 \text{ Median} = \text{Mode} + \text{Mean}$
 - d. $3 \text{ Median} = \text{Mode} - 2 \text{ Mean}$

Q5. The mean of following distribution is:

| | | | | |
|-------|----|----|----|----|
| x_i | 11 | 14 | 17 | 20 |
| f_i | 3 | 6 | 8 | 7 |

- a. 15.6 b. 17 c. 14.8 d. 16.4

Q6. Consider the following frequency distribution:

| | | | | | |
|-----------|-----|------|-------|-------|-------|
| Classes | 0-5 | 6-11 | 12-17 | 18-23 | 24-29 |
| Frequency | 13 | 10 | 15 | 8 | 11 |

The upper limit of the median class is:

- a. 17 b. 17.5 c. 18 d. 18.5

Q7. The class interval of a given observation is 10 to 15, then the class mark for this interval will be:

- a. 11.5 b. 12.5 c. 12 d. 14

Q8. Mode is the_____.

- a. middle most frequent value b. least frequent value
c. maximum frequent value d. none of these

Q9. For the following distribution:

| | | | | | |
|-----------|------|-------|-------|-------|-------|
| Classes | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 |
| Frequency | 13 | 10 | 15 | 8 | 11 |

The number of students who got marks less than 30 is:

- a. 13 b. 25 c. 10 d. 12

Q10. The mode and mean is given by 7 and 8, respectively. Then the median is:

- a. $1/13$ b. $13/3$ c. $23/3$ d. 33

Q11. What is the algebraic sum of the deviations of a frequency distribution from its mean?

Q12. Find the mean of first five prime numbers.

Q13. Is range a measure of central tendency in a distribution?

Q14. Write the formula for finding the mode of grouped data.

Q15. Find mean of the following frequency distribution :

| | | | | | |
|-----------|------|-------|-------|-------|--------|
| Classes | 0-20 | 20-40 | 40-60 | 60-80 | 80-100 |
| Frequency | 15 | 18 | 21 | 29 | 17 |

Q16. Find mode :

| | | | | | | | | |
|----------------|------|-------|-------|-------|-------|-------|-------|-------|
| Class Interval | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 | 50-60 | 60-70 | 70-80 |
| Frequency | 5 | 8 | 7 | 12 | 28 | 20 | 10 | 10 |

Q17. Calculate the mean for the following distribution:

| | | | | | |
|---|---|---|----|----|---|
| x | 5 | 6 | 7 | 8 | 9 |
| f | 4 | 8 | 14 | 11 | 3 |

Q18. Calculate the mode from the following data:

| | | | | | | | | |
|-----------|------|-------|-------|-------|-------|-------|-------|-------|
| Classes | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 | 50-60 | 60-70 | 70-80 |
| Frequency | 5 | 8 | 7 | 12 | 28 | 20 | 10 | 10 |

Q19. Calculate the median from the following data:

| | | | | | | | | |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|
| Classes | 15-25 | 25-35 | 35-45 | 45-55 | 55-65 | 65-75 | 75-85 | 85-95 |
| Frequency | 8 | 10 | 15 | 25 | 40 | 20 | 15 | 7 |

Q20. If the mean of the distribution is 54, find the value of p:

| | | | | | |
|-----------|------|-------|-------|-------|--------|
| Classes | 0-20 | 20-40 | 40-60 | 60-80 | 80-100 |
| Frequency | 7 | p | 10 | 9 | 13 |

Q21. The mean of the frequency table is 50. But the frequencies f_1 and f_2 are missing. Find them.

| | | | | | | |
|-----------|------|-------|-------|-------|--------|-------|
| Class | 0-20 | 20-40 | 40-60 | 60-80 | 80-100 | Total |
| Frequency | 17 | f_1 | 32 | f_2 | 19 | 120 |

Q22. If the median of the frequency distribution is 46, find the missing frequencies:

| | | | | | | | | |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|
| Variable | 10-20 | 20-30 | 30-40 | 40-50 | 50-60 | 60-70 | 70-80 | Total |
| Frequency | 12 | 30 | p | 65 | q | 25 | 18 | 229 |

Q23. The following table gives the daily income of 50 workers of a factory:

| | | | | | |
|-------------------|---------|---------|---------|---------|---------|
| Daily income | 100-120 | 120-140 | 140-160 | 160-180 | 180-200 |
| Number of workers | 12 | 14 | 8 | 6 | 10 |

Find the mean, mode and median of the above data.

Q24. Calculate the missing frequency from the following distribution, it being given that the median of the distribution is 24.

| | | | | | |
|--------------|------|-------|-------|-------|-------|
| Age in years | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 |
| No of houses | 5 | 25 | p | 18 | 7 |

Case-study based questions:

Q25. The COVID-19 pandemic, also known as corona virus pandemic, is an ongoing pandemic of corona virus disease caused by the transmission of severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) among humans.



The following tables shows the age distribution of case admitted during a day in two different hospitals

Table 1

| Age (in years) | 5-15 | 15-25 | 25-35 | 35-45 | 45-55 | 55-65 |
|-------------------|------|-------|-------|-------|-------|-------|
| No. of cases | 6 | 11 | 21 | 23 | 14 | 5 |

Table 2

| Age (in years) | 5-15 | 15-25 | 25-35 | 35-45 | 45-55 | 55-65 |
|-------------------|------|-------|-------|-------|-------|-------|
| No. of cases | 8 | 16 | 10 | 42 | 24 | 12 |

Refer to table 1

- i. The average age for which maximum cases occurred is:
 - a. 32.24
 - b. 34.36
 - c. 36.82
 - d. 42.24
- ii. The upper limit of modal class is:
 - a. 15
 - b. 25
 - c. 35
 - d. 45
- iii. The mean of the given data is:
 - a. 26.2
 - b. 32.4
 - c. 33.5
 - d. 35.4

Refer to table 2

- iv. The mode of the given data is:
 - a. 41.4
 - b. 48.2
 - c. 55.3
 - d. 64.6
- v. The median of the given data:
 - a. 32.7
 - b. 40.2
 - c. 42.3
 - d. 48.6

CHAPTER - 15 (PROBABILITY)

Solve the following questions:

- Q1.** The probability expressed as a percentage of a particular occurrence can never be :
- less than 100
 - less than 0
 - greater than 1
 - anything but a whole number
- Q2.** Which of the following is true?
- $0 \leq P(E) \leq 1$
 - $P(E) > 1$
 - $P(E) < 0$
 - $-\frac{1}{2} \leq P(E) \leq \frac{1}{2}$
- Q3.** The probability of passing a certain test is $\frac{x}{24}$. If the probability of not passing it is $\frac{7}{8}$, then x is equal to :
- 2
 - 3
 - 4
 - 6
- Q4.** A card is drawn from a deck of 52 cards. The event E is that card is not an ace of hearts. The number of outcomes favourable to E is _____.
- 4
 - 13
 - 48
 - 51
- Q5.** A letter is chosen at random from the English alphabet. Find the probability that the letter chosen succeeds X.
- $\frac{1}{13}$
 - $\frac{1}{52}$
 - $\frac{1}{26}$
 - $\frac{1}{2}$
- Q6.** Two different dice are tossed together. The probability that the product of the two numbers on the top of the dice is 6, is :
- $\frac{2}{3}$
 - $\frac{1}{3}$
 - $\frac{1}{9}$
 - $\frac{7}{36}$
- Q7.** Two coins are tossed simultaneously. The probability of getting exactly one head is _____.
- $\frac{1}{2}$
 - $\frac{1}{4}$
 - $\frac{1}{6}$
 - None of these
- Q8.** A bag contains 24 balls of which x are red, 2x are white and 3x are blue. A ball is selected at random. What is the probability that the drawn ball is white or blue?
- $\frac{1}{2}$
 - 2
 - $\frac{5}{6}$
 - $\frac{7}{12}$
- Q9.** A letter is chosen at random from the letters of the word 'ASSASSINATION', then the probability that the letter chosen is a vowel is in the form of $\frac{6}{2x+1}$, then x is equal to :
- 5
 - 6
 - 7
 - 8
- Q10.** Two unbiased coins are tossed simultaneously. Find the probability of getting
- one head
 - one tail
 - two heads
 - at least one head
 - at most one head.
- Q11.** If the probability of answering a question correctly is 0.3, what is the probability of getting it wrong?

- Q12.** A game of chance consists of spinning an arrow which comes to rest pointing at one of the numbers marked from 1 to 10. What is the probability that it will point at an even number?
- Q13.** If three coins are tossed simultaneously, then what is the probability of getting at least one head and one tail?
- Q14.** Tickets numbered 1 to 20 are mixed up and then a ticket is drawn at random. What is the probability that the ticket drawn bears a number which is a multiple of 3?
- Q15.** What is the probability that a leap year selected at random will contain 53 Sundays and 53 Mondays?
- Q16.** A number is chosen at random from the numbers -3, -2, -1, 0, 1, 2, 3. What will be the probability that square of this number is less than or equal to 1?
- Q17.** A carton of 24 bulbs contain 6 defective bulbs. One bulb is drawn at random. What is the probability that the bulb is not defective? If the bulb selected is defective and it is not replaced and a second bulb is selected at random from the rest, what is the probability that the second bulb is defective?
- Q18.** A child's game has 8 triangles of which 3 are blue and rest are red, and 10 squares of which 6 are blue and rest are red. One piece is lost at random. Find the probability that is a
- triangle,
 - square,
 - square of the blue colour,
 - triangle of the red colour.
- Q19.** The King, Queen and Jack of clubs are removed from a pack of 52 cards and then the remaining cards are well shuffled. A card is selected from the remaining cards. Find the probability of getting a card
- of spade,
 - of black king,
 - of club,
 - of jacks.
- Q20.** The probability of selecting a blue marble at random from a jar that contains only blue, black and green marbles is $\frac{1}{5}$. The probability of selecting a black marble at random from the same jar is $\frac{1}{4}$. If the jar contains 11 green marbles, find the total number of marbles in the jar.
- Q21.** An integer is chosen at random between 1 and 100. Find the probability that it is
- divisible by 8,
 - not divisible by 8.
- Q22.** All the three face cards of spades are removed from a pack of 52 cards. A card is drawn at random from the remaining cards, find the probability of getting
- a black face card,
 - neither a king nor a red card,
 - either black or queen.
- Q23.** Cards are numbered from 7 to 51, one card is drawn at random, find the probability of getting
- prime number less than 20,
 - not divisible by 10,
 - divisible by 2 or 3.

Case-study based questions:

Q24. In a club, men are playing the card game. A man named Sanjeev draw a card from a well shuffled deck of cards.



Based on the above information, give the answer of the following questions :

i. Find the probability of getting a king of black colour :

- a. $\frac{1}{26}$ b. $\frac{1}{12}$ c. $\frac{1}{52}$ d. $\frac{1}{4}$

ii. Find the probability of getting a face card:

- a. $\frac{1}{26}$ b. $\frac{1}{13}$ c. $\frac{2}{13}$ d. $\frac{3}{13}$

iii. Find the probability of getting a jack of clubs:

- a. $\frac{1}{26}$ b. $\frac{1}{52}$ c. $\frac{3}{52}$ d. $\frac{3}{26}$

iv. Find the probability of getting a black face card:

- a. $\frac{3}{13}$ b. $\frac{1}{13}$ c. $\frac{1}{52}$ d. $\frac{3}{26}$

v. Find the probability of getting a diamond:

- a. $\frac{1}{26}$ b. $\frac{1}{13}$ c. $\frac{1}{52}$ d. $\frac{1}{4}$